

TRYPANOCIDES AND  
ANTI-PROTOZOALS

LARGE  
RUMINANTS



# ACTIDIUM

**TREAT AND PREVENT  
TRYPANOSOMIASIS...**



**...WITH A HOMIDIUM  
CHLORIDE-BASED  
TRYPANOCIDE**





LARGE  
RUMINANTS

TRYPANOCIDES AND  
ANTI-PROTOZOALS

# ACTIDIUM



## INDICATIONS

Treatment

Prevention



**Trypanosomes**

(*Trypanosoma vivax*, *T. brucei* and *T. congolense*)



## COMPOSITION

1 Tablet



**HOMIDIUM CHLORIDE** .....250mg

**PHENAZONE** .....250mg

Excipients ..... q.s. to 1 tablet



## ADMINISTRATION



## PHARMACEUTICAL FORM

Injectable  
Route



Deep Intramuscular  
Injection in the  
neck

Tablet for Injectable Solution



## DOSAGE

### Dilution

Dissolve 1 tablet in 10 mL of sterile water.

### Administration

2 mL per 50 kg BW  
4 mL per 100 kg BW  
8 mL per 200 kg BW  
10 mL per 250 kg BW



Number of animals treated per tablet:  
1 animal weighing 250 kg BW

**Duration of use**  
**Duration of action**



**Single dose**  
**6 weeks**



## WITHDRAWAL PERIOD



**MILK**  
**0 DAY**



**MEAT AND OFFAL**  
**28 DAYS**



## PRESENTATIONS



- Tablet for  
Injectable Solution
- 20-Tablet Vial
- Box of 10 vials

