

**EXTERNAL
ANTIPARASITICS**

ALPHALOBS

**RUMINANTS
POULTRY - SWINE
EQUINES**



***TREAT ALL EXTERNAL PARASITES
IN RUMINANTS, SWINE
AND POULTRY...***



***...WITH A REFERENCE
ALPHACYPERMETHRIN-BASED
SOLUTION***





RUMINANTS
POULTRY - SWINE
EQUINES

EXTERNAL
ANTIPARASITICS

ALPHALOBS



INDICATIONS



Mange



Ticks



Lice



Flies



COMPOSITION

100 mL of solution

ALPHACYPERMETHRINE 10% m/v

Excipients q.s. to 100 mL



ADMINISTRATION



PHARMACEUTICAL FORM

Solution for Dipping or Spraying

Cutaneous Use



Dipping



Local Application
by Spraying



+



=



+



=



DOSAGE



100 mL

+



200L

=



First Dip

100 mL

125L

Subsequent Dips



10mL

+



20L

=



3 L of diluted solution per animal

Duration of Use



Single Dose

Administration Frequency



Ticks

- Ticks: In cases of heavy infestation, spray every week. A year-round control programme is recommended.



Flies

- Tsetse Flies: Apply for 2 to 4 weeks, depending on the level of infestation.



Mange

- Mange: 2 treatments at 10-day intervals



Lice

- Lice: 1 treatment as needed.

- Flies and other harmful insects: 1 treatment every 6 to 8 days, or according to the level of infestation.



WITHDRAWAL PERIODS



MILK
5 MILKINGS



MEAT AND OFFAL
8 DAYS



PRESENTATIONS



Solution



Cutaneous



100 mL Vial

X1



Solution



Cutaneous



20 mL Vial



Box of 25 Vials

X25



Solution



Cutaneous



1 L Vial

X1

