

**INTERNAL
ANTIPARASITICS**

RUMINANTS



DISTOCUR

TREAT ALL WORMS IN RUMINANTS...



***...WITH A BITHIONOL SULFOXIDE-BASED
INTERNAL ANTIPARASITIC
IN BOLUS FORM***



DISTOCUR



INDICATIONS

• Treatment • Prevention •



Tapeworms

Thysaniezia ovilla, Moniezia, Stilesia



Roundworms

Oesophagostomum



Liver Flukes

Fasciola Hepatica



Paramphistomes



COMPOSITION

One bolus 

BITHIONOL SULFOXIDE2g

Excipients q.s. to 2,7g



ADMINISTRATION



PHARMACEUTICAL FORM

Oral Route



Direct Oral
Administration

Bolus



DOSAGE



1 bolus per 50 kg BW



1/2 bolus per 25 kg BW

Duration of Use



Single dose

Administration schedule

- At the beginning of the rainy season, followed by a booster dose 15 days later
- Before vaccination, at weaning, before parturition and before mating



WITHDRAWAL PERIODS



MILK
0 DAY



MEAT AND OFFAL
15 DAYS



PRESENTATIONS



 Bolus
 Oral
 100-Boli Jar



 Bolus
 Oral
 1000-Boli Jar

