

ANTI-INFECTIVES

**RUMINANTS
POULTRY - SWINE**



DOXYLOBS 500

**TREAT INFECTIONS CAUSED BY
DOXYCYCLINE-SENSITIVE BACTERIA...**



**...WITH AN ORAL POWDER
FORMULATED WITH A SINGLE
EFFECTIVELY DOSED ANTIBIOTIC**



DOXYLOBS 500



INDICATIONS



Bacteria

Doxycycline-sensitive bacteriae



COMPOSITION

One packet of 100g

DOXYCYCLINE (as hydrochloride) 50g

Excipients q.s. to 100g



ADMINISTRATION



PHARMACEUTICAL FORM


Oral Route



Drinking Water



Feed



Pellets for Oral Solution

Poudre for Medicated Feed



DOSAGE



100 g per 500–1000 L of drinking water

Duration of Use



Pendant 3 à 5 jours



1 g per 100 kg BW, diluted in a small amount of water or mixed with feed **twice daily**

Duration of Use



Pendant 3 à 5 jours


- For use only in pre-ruminant calves, lambs and kids
- Do not use in adult ruminants



WITHDRAWAL PERIODS



MEAT AND OFFAL



7 DAYS

14 DAYS

8 DAYS

- Do not use in dairy cows producing milk for human consumption or in laying hens producing eggs for human consumption.



PRESENTATIONS



- Powder
- Oral solution or feed
- 100 g sachet
- Box of 25 sachets



- Powder
- Oral solution or feed
- 1 kg jar

