



LIVACOX Q

**VACCINATE LAYING
HENS AGAINST COCCIDIOSIS...**



**...WITH AN EFFECTIVE
AND EASY-TO-USE
LIVE VACCINE**





POULTRY

VACCINES

LIVACOX Q



INDICATIONS

Prevention



Coccidia

Live attenuated quadrivalent vaccine against coccidiosis in domestic poultry (*Eimeria acervulina*, *Eimeria maxima*, *Eimeria tenella*, *Eimeria necatrix*)



COMPOSITION

Vial Contains 1,000 or 5,000 Vaccine Doses

One Vaccine Dose

- **COCCIDIAL OOCYSTS** *Eimeria acervulina* 3 -5. 10², *Eimeria tenella* 3 -5. 10², *Eimeria maxima* 3 -5. 10², *Eimeria necatrix* 1. 10²
- Excipients : q.s. to 1 dose



ADMINISTRATION



PHARMACEUTICAL FORM

Oral / Spray Solution to be Diluted

Oral Route



Drinking Water

Respiratory Route



Spray Administration



DOSAGE

10 mL of preparation = 1,000 doses

50 mL of preparation = 5,000 doses

Dilution

Dilute in drinking water.

1 L of drinking water / day of age / 1,000 birds

Example: For 1,000 birds 4 days of age

→ Dilute 10 mL in 4 L of drinking water.



For 1-Day-Old Chicks

Dilute in drinking water

1 L of drinking water / 1,000 birds

For 1,000 Birds → Dilute 10 mL in 1 L of drinking water.



Administration

- The birds must consume the preparation within 2 hours (Consumption may be encouraged by withholding drinking water for 2 hours before vaccination).

- The vaccine may be diluted in water containing food colouring (to facilitate visual monitoring of application). It is then sprayed onto the chicks. The chicks must remain in the crate for at least 3 hours.

Duration of Use



Single Dose

IMMUNITY: develops 10 days after vaccination and lasts throughout the bird's lifetime.

- Vaccinate healthy birds only.
- The vaccine should be administered to chicks between 1 and 10 days of age; however, hens may be vaccinated at any age during the rearing period.
- Feed given to the birds from 2 days before until 14 days after vaccination must not contain anticoccidials at therapeutic doses or sulfonamides.



WITHDRAWAL PERIODS



MEAT AND OFFAL
0 DAY



MILK
0 DAY



PRESENTATIONS



- Solution
- Drinkable
- Nebulizable
- 10 mL Vial – 1,000 Doses
- Box of 10 Vials



- Solution
- Drinkable
- Nebulizable
- 50 mL Vial – 5,000 Doses
- Box of 10 Vials

STORAGE

- Store at 2–8°C
- Shelf life: 9 months

