

VACCINES

RUMINANTS
EQUINES



LOBALIVE ANTHRAX

**VACCINATE RUMINANTS
AND HORSES AGAINST
ANTHRAX...**



**...WITH AN EFFECTIVE
AND EASY-TO-USE
LIVE VACCINE**





RUMINANTS
EQUINES

VACCINES

LOBALIVE ANTHRAX



INDICATIONS

Prevention



Bacteria

Live attenuated vaccine against anthrax (*Bacillus anthracis*).



COMPOSITION

Lyophilized Vial Contains 50 or 100 Vaccine Doses
One Vaccine Dose Δ

- BACILLUS ANTHRACIS**, Sterne 34 F2 strain (adjuvanted with saponin) — 10^7 viable spores
Saponin concentration: 1–2% (the spores are lyophilized with saponin)
- Excipient : q.s. to 1 dose



ADMINISTRATION



PHARMACEUTICAL FORM

Injectable Route



- Subcutaneous
- Loose skin of the neck
 - Loose skin of the neck or inner thigh
 - Inner thigh
 - Skin of the neck



DOSAGE

Dilution

Reconstitute the vaccine:

- 1 vial of 100-dose lyophilisate with 100 mL of diluent.
- 1 vial of 50-dose lyophilisate with 50 mL of diluent

Administration

- : 1 mL
- : 0.5 mL
- : 0.5 mL
- : 1 mL

Duration of Use
Duration of Action



Single Dose
Booster: Annually

IMMUNITY: Immunity develops 10 days after vaccination and lasts for 12 months.



WITHDRAWAL PERIOD



MEAT AND OFFAL
6 WEEKS

DO NOT ADMINISTER TO
PREGNANT FEMALES



PRESENTATIONS



X1

- Lyophilisate for Injectable Solution
- Subcutaneous (after reconstitution with diluent)
- Lyophilized Vial to be reconstituted with 50 mL of diluent (50 doses)



X1

- Lyophilisate for Injectable Solution
- Subcutaneous (after reconstitution with diluent)
- Lyophilized Vial to be reconstituted with 100 mL of diluent (100 doses)

STORAGE

- Store at 4°C
- Shelf life: 2 years

