

VACCINES

RUMINANTS



LOBALIVE[®] DNCB

***VACCINATE RUMINANTS AGAINST
LUMPY SKIN DISEASE (LSD)...***



***...WITH A LIVE ATTENUATED LYOPHILIZED
VACCINE PROVIDING RAPID AND
LONG-LASTING IMMUNITY***





LOBALIVE[®]

DNCB



INDICATIONS

Prevention



Virus

Live attenuated lyophilized vaccine against Lumpy Skin Disease (LSD)
(Capripoxvirus)



COMPOSITION

Each vial contains lyophilized vaccine containing 50 doses

1 mL of reconstituted lyophilisate contains

- **10⁴ TCID₅₀** of live attenuated lyophilized virus
- Vaccine strain: Capripoxvirus strain PANVAC



ADMINISTRATION



PHARMACEUTICAL FORM

Injectable
Route



Subcutaneous
Injection
Inner thigh



DOSAGE

Dilution

Reconstitute the vaccine as follows: 1 vial containing 50 doses of lyophilisate + 50 mL of diluent

Administration

Dose: 1 mL per bovine
Only healthy animals should be vaccinated.

Duration of Use
Duration of Action



Single dose
Booster vaccination every 6 months

IMMUNITY : 8 days after vaccination and lasting for 2 years.

- Maternal antibodies in calves may influence the effectiveness of the vaccine.
 - Calves born to vaccinated cows should be vaccinated at 6 months of age.
 - Calves born to unvaccinated cows may be vaccinated at any age.
- Although this vaccine has been extensively tested under a wide range of conditions, lack of efficacy may occur for various reasons. If this is suspected, consult a veterinarian.



WITHDRAWAL PERIOD



MEAT
21 DAYS

* The 21-day withdrawal period is a standard withdrawal period established under South African legislation for all vaccine products. If required by national regulations, an alternative withdrawal period may also be authorized.



PRESENTATIONS



X1



Lyophilized powder for injectable solution

Subcutaneous administration (after reconstitution with the diluent)

One vial of lyophilized vaccine to be reconstituted with 50 mL of diluent (50 doses)

STORAGE

- +4 and +8°C
- Shelf life: 2 years

