



LOBALIVE NEW IB

***PROTECT POULTRY AGAINST
NEWCASTLE DISEASE AND
INFECTIOUS BRONCHITIS...***



***...WITH AN EFFECTIVE AND
EASY-TO-USE LIVE VACCINE***





POULTRY

VACCINES

LOBALIVE NEW IB



INDICATIONS

Prevention



Virus

Live attenuated vaccine against Newcastle Disease (Paramyxovirus) and Infectious Bronchitis (Coronavirus)



COMPOSITION

Lyophilized Vial Contains 500–1000 Vaccine Doses
One vaccine dose

- **AVIAN INFECTIOUS BRONCHITIS VIRUS** (souche H120) Minimum $10^{3.8}$ EID₅₀*
- **NEWCASTLE DISEASE VIRUS** (souche La Sota) Minimum $10^{6.5}$ EID₅₀*
- Excipients : q.s. to 1 dose

* EID₅₀ = Embryo Infectious Dose 50%: viral titre capable of infecting 50% of inoculated embryos.



ADMINISTRATION

Oral Route



Drinking
Water

Oculonasal Route



Instillation

Respiratory Route



Nebulization



PHARMACEUTICAL FORM

Lyophilisate for

Oral / Oculonasal / Spray Suspension



* Diluent sold separately.



DOSAGE

Dilution

For 500 birds aged 5–10 days, dissolve 1000 vaccine doses in 5–10 L of drinking water (5–10 mL per bird).

For 500 birds over 10 days of age, dissolve 1000 vaccine doses in 10–20 L of drinking water (10–20 mL per bird).

Dissolve 1000 vaccine doses in 30 mL of water and administer one drop into one eye or one nostril of each bird.

For a flock of 1000 birds, dissolve 1000 vaccine doses in 10–40 L of water and spray over the birds.

Administration

- Dissolve the required amount of vaccine in fresh, non-chlorinated water, free from detergents, disinfectants and metallic ions.
- Withdraw drinking water 4 hours before vaccination to ensure rapid consumption of the vaccine.
- The reconstituted vaccine must be consumed within 4 hours.
- Reconstitute the vaccine according to the instructions and administer one drop (0.03 mL) into one eye or one nostril using a suitable dropper.
- Hold the bird's head slightly tilted and allow the vaccine to spread over the entire surface of the eye before releasing the bird.
- Use suitable spraying equipment to distribute the vaccine evenly throughout the poultry house.

Duration of Use



Single Dose

- The vaccine may be used for primary vaccination or booster vaccination in broilers, pullets, adult laying hens and chicks from 5–10 days of age onwards.



WITHDRAWAL PERIODS



MEAT AND OFFAL
0 DAY



EGGS
0 DAY



PRESENTATIONS



X1

- Lyophilisat solution
- Orale
- Oculonasale
- Spray Suspension
- 500–1000-Dose Lyophilized Vial

STORAGE

- 2–8°C
- Shelf life: 2 years.

