



LOBALIVE NEW

***PROTECT POULTRY AGAINST
NEWCASTLE DISEASE...***



***...WITH AN EFFECTIVE AND
EASY-TO-USE LIVE VACCINE***





POULTRY

VACCINES

LOBALIVE NEW



INDICATIONS

Prevention



Virus

Live attenuated Newcastle disease vaccine (Paramyxovirus)



COMPOSITION

Lyophilized Vial Contains 500–1000 Vaccine Doses

One vaccine dose

- **NEWCASTLE DISEASE VIRUS** (La Sota strain) Minimum $10^{6.5}$ EID₅₀*
- Excipients : q.s. to 1 dose

* EID₅₀ = Embryo Infectious Dose 50%: viral titre capable of infecting 50% of inoculated embryos.



ADMINISTRATION

Oral Route



Drinking Water

Oculonasal Route



Instillation

Respiratory Route



Nebulization



PHARMACEUTICAL FORM

Lyophilisate for Oral, Oculonasal or Spray Suspension



DOSAGE

Dilution

For 500 birds aged 5–10 days, dissolve 1000 vaccine doses in 5–10 L of drinking water (5–10 mL per bird).

For 500 birds over 10 days of age, dissolve 1000 vaccine doses in 10–20 L of drinking water (10–20 mL per bird).

Dissolve 1000 vaccine doses in 30 mL of water and administer one drop into one eye or one nostril of each bird.

For a flock of 1000 birds, dissolve 1000 vaccine doses in 10–40 L of water and spray over the birds.

Administration

- Dissolve the required amount of vaccine in fresh, non-chlorinated water, free from detergents, disinfectants and metallic ions.
- Withdraw drinking water 4 hours before vaccination to ensure rapid consumption of the vaccine.
- The reconstituted vaccine must be consumed within 4 hours.

- Reconstitute the vaccine according to the instructions and administer one drop (0.03 mL) into one eye or one nostril using a suitable dropper.
- Hold the bird's head slightly tilted and allow the vaccine to spread over the entire surface of the eye before releasing the bird.

- Use suitable spraying equipment to distribute the vaccine evenly throughout the poultry house.

Duration of Use



Single Dose

- The vaccine may be used for primary vaccination or booster vaccination in broilers, pullets, adult laying hens and chicks from 5–10 days of age onwards.



WITHDRAWAL PERIODS



**MEAT AND OFFAL
0 DAY**



**EGGS
0 DAY**



PRESENTATIONS



- Lyophilisate solution
- Oral
- Oculonasal
- Spray Suspension
- 500–1000-Dose Lyophilized Vial

STORAGE

- 2–8°C
- Shelf life: 2 years.

