



LOBALIVE S&G POX

***VACCINATE AGAINST
SHEEP POX AND GOAT POX...***



***...WITH AN EFFECTIVE
AND EASY-TO-USE LIVE VACCINE***





LOBALIVE S&G POX



INDICATIONS

Prevention



Virus

Live attenuated vaccine against sheep pox and goat pox (Poxvirus)



COMPOSITION

Lyophilized Vial Contains 50 or 100 Vaccine Doses
One Vaccine Dose Δ

- SHEEP POX AND GOAT POX VIRUS**, CVCC AV41 strain — titre $\geq 1 \times 10^{3.5}$ TCID₅₀.
TCID₅₀: 50% Tissue Culture Infectious Dose
- Excipient : q.s. to 1 dose



ADMINISTRATION



PHARMACEUTICAL FORM

Injectable
Route



Subcutaneous
Injection site: Inner thigh
or under the tail



DOSAGE

Dilution

Reconstitute the vaccine:

- 1 vial of 100-dose lyophilisate with 100 mL of diluent
- 1 vial of 50-dose lyophilisate with 50 mL of diluent

Administration

Dose: 1 mL per animal
Only healthy animals should be vaccinated.

Duration of Use
Duration of Action



Single Dose
Booster: Annually

IMMUNITY: develops 4 to 5 days after vaccination and lasts for 12 months.



WITHDRAWAL PERIODS



MEAT AND OFFAL
0 DAY



MILK
0 DAY



PRESENTATIONS



X1

- Lyophilisate for Injectable Solution
- Subcutaneous (after reconstitution with diluent)
- Lyophilized Vial to be reconstituted with 50 mL of diluent (50 doses)

STORAGE

- 24 months below -15°C
- 18 months at $2-8^{\circ}\text{C}$



X1

- Lyophilisate for Injectable Solution
- Subcutaneous (after reconstitution with diluent)
- Lyophilized Vial to be reconstituted with 100 mL of diluent (100 doses)

