

TRYPANOCIDES AND
ANTI-PROTOZOALS

RUMINANTS



LOBAZEN

**TREAT TRYPANOSOMIASIS, BABESIOSIS
AND THEILERIOSIS...**



**...WITH PROVEN TRYPANOCIDAL
ACTIVITY**



LOBAZEN



INDICATIONS

Treatment



Trypanosomiasis

(*Trypanosoma congolense*,
T. vivax et *T. Brucei*.)



Theileriosis

(*Theileria*
annulata)



Babesiosis

(*Babesia bovis*, *B. bigemina*, *B. ovis*, and other *Babesia* species.)



COMPOSITION

One packet of 2.36 g 

DIMINAZENE Diaceturate 1.05g
Antipyrine 1.31g

One packet of 23.6 g 

DIMINAZENE Diaceturate 10.5g
Antipyrine 13.1g



ADMINISTRATION

Injectable
Route



Deep
intramuscular
injection



PHARMACEUTICAL FORM

Pellets for injectable solution



DOSAGE

Dilution

-  2.36 g (1 sachet) for 15 mL of water
-  23.6 mg (1 sachet) for 150 mL of water

Administration

1 mL per 20 kg BW
5 mL per 100 kg BW
10 mL per 200 kg BW
15 mL per 300 kg BW



Duration of use



Single dose



WITHDRAWAL PERIOD



MILK
5 DAYS



MEAT AND OFFAL
21 DAYS



PRESENTATIONS



Powder for
injectable solution
 2.36 g sachet
 Box of 10 sachets



Powder for
injectable solution
 23.6 g sachet
 Box of 10 sachets

