



LOBIVAC 8 DHPPiL

**VACCINATE DOGS AGAINST
CANINE DISTEMPER (D), RUBARTH'S
HEPATITIS (H), PARVOVIROSIS (P),
KENNEL COUGH (PI) AND
LEPTOSPIROSIS (L)...**



**...WITH AN EFFECTIVE
AND EASY-TO-USE VACCINE**





DOGS

VACCINES

LOBIVAC 8 DHPPiL



Prevention



Virus



Bacteria

Vaccine against canine distemper (D), infectious canine hepatitis – Rubarth's hepatitis (H), parvovirus (P), kennel cough – parainfluenza (Pi), and leptospirosis (L).



COMPOSITION

1 Lyophilisate Vial + 1 Diluent Vial Contains 1 Vaccine Dose
One Vaccin Dose (1mL)

Lyophilisate:

- **VIRUS FEBRIS CONTAGIOSAE CANIS ATTENUATUM** $10^{2.7} - 10^{4.5}$ EID₅₀,
- **VIRUS LARYNGOTRACHEITIDIS CONTAGIOSAE CANIS ATTENUATUM** $10^{3.5} - 10^{7.5}$ CCID₅₀,
- **PARVOVIRUS ENTERITIDIS CANIS ATTENUATUM** $10^{2.1} - 10^{3.6}$ HAU,
- **VIRUS PARAINFLUENSIS CANIS ATTENUATUM** $10^{3.0} - 10^{6.0}$ CCID₅₀,
- **LEPTOSPIRA ICTEROHAEMORRHAGIAE INACTIVATA, LEPTOSPIRA GRIPPOTYPHOSA INACTIVATA, LEPTOSPIRA SEJROE INACTIVATA** q.s. to obtain an experimental serological response in animals with a minimum specific antibody titre of 1:32.

EID₅₀: 50% Embryo Infective Dose

CCID₅₀: 50% Cell Culture Infective Dose.

HAU: Haemagglutination Unit

Diluent:

- Excipient: Water for Injection — 1 mL



ADMINISTRATION

Injectable
Route



Subcutaneous



PHARMACEUTICAL FORM

Lyophilisate and Diluent for
Injectable Solution



Lyophilisate

Diluent



DOSAGE

Dilution

Dissolve the lyophilisate in 1 mL of diluent immediately before use.

Administration

Dose: 1 mL per animal
Only healthy animals should be vaccinated from 6 weeks of age.

Duration of Use
Duration of Action



Primary Vaccination

(One injection at 6–8 weeks of age, followed by a second injection 2–3 weeks later)*

Booster: Annually

IMMUNITY: develops 2 to 3 weeks after primary vaccination and lasts for 1 year.

*A third vaccine dose is recommended 3–4 weeks after the second dose in young animals with specific colostral antibodies and in young animals that received the second dose before 12 weeks of age.



WITHDRAWAL PERIOD

Do not use in animals intended for human consumption.



PRESENTATIONS



Lyophilisate + Diluent for Mixing

Subcutaneous Injection

1 Lyophilisate Vial

+ 1 mL Diluent Vial

Box of 6 Lyophilisate Vials

+ 6 Diluent Vials (Each Box Contains 6 Doses)

STORAGE

- Store at 2–8°C
- Shelf life: 2 years

