



***VACCINATE LAYERS AND CHICKENS
AGAINST AVIAN INFLUENZA,
NEWCASTLE DISEASE AND
INFECTIOUS BRONCHITIS...***



**...WITH AN EFFECTIVE AND
EASY-TO-USE COMBINED
INACTIVATED VACCINE**





POULTRY

VACCINES

LOBIVAC

AI-NEW-IB-EDS



INDICATIONS

Prevention



Virus

Inactivated vaccine indicated for the prevention of Avian Influenza (Influenzavirus), Newcastle Disease (Paramyxovirus), Infectious Bronchitis (Coronavirus) and Egg Drop Syndrome (EDS, Adenovirus).



COMPOSITION

The inactivated vaccine contains:

- **AVIAN INFLUENZA VIRUS** (H9 subtype, NJ02 strain), minimum viral titre prior to inactivation $\geq 4 \times 10^{8.0} \text{ DIE}_{50} / 0.1 \text{ mL}^*$
- **NEWCASTLE DISEASE VIRUS** (LaSota strain), minimum viral titre prior to inactivation $\geq 4 \times 10^{8.0} \text{ DIE}_{50} / 0.1 \text{ mL}^*$
- **AVIAN INFECTIOUS BRONCHITIS VIRUS** (M41 strain), minimum viral titre prior to inactivation $\geq 4 \times 10^{6.7} \text{ DIE}_{50} / 0.1 \text{ mL}^*$
- **EGG DROP SYNDROME VIRUS** (EDS) (AV127 strain), with a hemagglutination inhibition titre prior to inactivation $\geq 1:10,240$;

* DIE_{50} = dose infectant 50 % des embryons : titre viral causant une infection chez 50 % des embryons inoculés avec le virus

**U.IH = Unité Inhibition de l'hémagglutination



ADMINISTRATION




PHARMACEUTICAL FORM

Injectable
Route



Subcutaneous Injection at
the back of the neck
Intramuscular injection in
the anterior portion of the
breast muscle

Injectable Suspension



DOSAGE

Dilution

Use the product as supplied.

Administration

- 0.3 mL per bird younger than 3 weeks.
- 0.5 mL per bird older than 3 weeks.

Duration of Use
Duration of Action



Single dose

**Booster vaccination: 3 months
after vaccination for chicks that
received a 0.3 mL dose.**

**6 months after vaccination for
birds that received a 0.5 mL dose**

- 500 mL vial: 1000 doses (birds older than 3 weeks) or 1665 doses (chicks).



WITHDRAWAL PERIOD



MEAT AND OFFAL
21 DAYS



EGGS
42 DAYS



PRESENTATIONS



X1

- Injectable solution
- Subcutaneous or intramuscular administration
- 500 mL vial

STORAGE CONDITIONS

- Shelf life: 12 months when stored at 2–8°C

