



LOBIVAC AI-NEW-IB

**VACCINATE CHICKENS AGAINST
AVIAN INFLUENZA, NEWCASTLE DISEASE
AND INFECTIOUS BRONCHITIS...**



**...WITH AN EFFECTIVE AND
EASY-TO-USE COMBINED
INACTIVATED VACCINE**





POULTRY

VACCINES

LOBIVAC

AI-NEW-IB



INDICATIONS

Prevention



Virus

Inactivated vaccine against avian influenza (Influenzavirus), Newcastle disease (Paramyxovirus) and infectious bronchitis (Coronavirus)



COMPOSITION

The inactivated vaccine contains:

- **AVIAN INFLUENZA VIRUS** H9N2 subtype, L strain), minimum viral titre prior to inactivation $10^{7.5} \text{ EID}_{50}/0.1\text{mL}^*$
- **NEWCASTLE DISEASE VIRUS** (LaSota strain), minimum viral titre prior to inactivation $10^{8.5} \text{ EID}_{50}/0.1\text{mL}^*$
- **AVIAN INFECTIOUS BRONCHITIS VIRUS** (M41 strain), minimum viral titre prior to inactivation $10^{7.0} \text{ EID}_{50}/0.1\text{mL}^*$

* EID_{50} = egg infective dose 50%: the viral titre required to infect 50% of inoculated embryonated eggs



ADMINISTRATION

Injectable
Route



Subcutaneous Injection at the back of the neck
Intramuscular injection in the anterior portion of the breast muscle



PHARMACEUTICAL FORM

Injectable Suspension



DOSAGE

Dilution

Use the product as supplied.

Administration

- 0.3 mL per bird younger than 3 weeks.
- 0.5 mL per bird older than 3 weeks.

Duration of Use
Duration of Action



Single dose
Booster vaccination: 3 months after vaccination for chicks that received a 0.3 mL dose.
6 months after vaccination for birds that received a 0.5 mL dose

- 500 mL vial: 1000 doses (birds older than 3 weeks) or 1665 doses (chicks).



WITHDRAWAL PERIOD



MEAT AND OFFAL
21 DAYS



EGGS
42 DAYS



PRESENTATIONS



x1

- Injectable solution
- Subcutaneous or intramuscular administration
- 500 mL vial

STORAGE CONDITIONS

- Shelf life: 12 months when stored at 2–8°C

