

VACCINES

LOBIVAC RAGE

DOGS - CATS - RUMINANTS
SWINE - EQUINES



**VACCINATE DOGS, CATS
AND OTHER ANIMAL SPECIES
AGAINST RABIES...**



**...WITH AN EFFECTIVE AND
EASY-TO-USE INACTIVATED
VACCINE**



LOBIVAC RAGE



INDICATIONS

Prevention



Virus

Inactivated Rabies Vaccine
(Rhabdoviridae)



COMPOSITION

Each vial contains one vaccine dose

One vaccine dose (1 mL)

- **INACTIVATED RABIES VIRUS MIN. 2 UI**
- Adjuvant: aluminum hydroxide gel 1.4-6.5mg
- Excipients :
 - Thiomersal < 0.1 mg
 - Other excipientsq.s. to 1mL



ADMINISTRATION

Injectable
Route



Subcutaneous injection



Intramuscular



PHARMACEUTICAL FORM

Injectable Suspension



DOSAGE

Dilution

Use the product as supplied (no dilution required).

Administration

Dose: 1 mL per animal
Only healthy animals from 12 weeks of age should be vaccinated.

Duration of Use
Duration of Action



Single dose (Primary Vaccination)*
Annual booster vaccination

IMMUNITÉ : Protective immunity develops 2–3 weeks after vaccination and lasts for 12 months.

•* In situations where there is a high risk of infection, animals may be vaccinated from 6 weeks of age. However, it is strongly recommended that a second dose be administered 2 to 4 weeks later.



WITHDRAWAL PERIOD



MEAT AND OFFAL
0 DAY



MILK
0 JOUR



PRESENTATIONS



X12

- Solution
- Injectable for subcutaneous or intramuscular administration
- 1 mL vial
- Box containing 12 vials (Each box contains 12 doses)

CONSERVATION

- Store at 2°C to 8°C
- Shelf life: 2 years

