

ANTI-INFECTIVES

LOBSPRAY

**RUMINANTS - SWINE
EQUINES - DOGS - CATS**



***TREAT SKIN WOUNDS
LOCALLY...***



***...WITH AN OXYTETRACYCLINE-BASED
ANTI-INFECTIVE SPRAY***





RUMINANTS - SWINE
EQUINES - DOGS - CATS

ANTI-INFECTIVES

LOBSPRAY



INDICATIONS

Treatment



Bacteria

Oxytetracycline-sensitive bacteria



COMPOSITION

210 mL of solution 

OXYTETRACYCLINE (as hydrochloride) 4,2g

Gentian Violet..... 1,05g

Excipients.....q.s. to 210mL



ADMINISTRATION



PHARMACEUTICAL FORM

Topical
Application



Spray onto the
affected or injured
area

Topical Spray Solution



DOSAGE



Spray for 1 to 2 seconds from a distance of approximately 20 cm (8 inches) from the affected or injured area while holding the container upright.

Duration of Use
Duration of Action



1 to 2 applications daily
For 3 days
(Do not exceed 10 days)



WITHDRAWAL PERIODS



MILK
0 DAY



MEAT AND OFFAL
0 DAY



PRESENTATIONS



x1



Spray Solution



Topical



Spray of 210mL

