

**INTERNAL
ANTIPARASITICS**

RUMINANTS



LOXINIL²⁵

***TREAT ROUNDWORMS,
LIVER FLUKES AND
BOT FLY LARVAE...***



***...WITH A NITROXYNIL-BASED
INJECTABLE INTERNAL
ANTIPARASITIC***



LOXINIL²⁵



INDICATIONS



Roundworms

Haemonchus contortus, *d'Haemonchus placei*,
Oesphagostomum radiatum, *Bunostomum*
phlebotomum, *Parafilaria Bovicola*



Liver Flukes

Fasciola hepatica (larval and
adult stages)



Bot Fly Larvae

Oestrosis



COMPOSITION

100 mL of solution

NITROXINIL 25g

Excipients q.s. to 100 mL



ADMINISTRATION

Injectable
Route




Injectable Solution



PHARMACEUTICAL FORM

Injectable Solution



DOSAGE



Subacute or
Chronic Infestations

1 mL per 25 kg BW



1,3 mL per 25 kg BW

Duration of Use



Single dose

Administration Schedule

- At the beginning of the rainy season, with a booster treatment 15 days later
- Before vaccination, at weaning, before parturition, before breeding...



WITHDRAWAL PERIODS



**DO NOT USE IN FEMALES PRODUCING MILK
FOR HUMAN CONSUMPTION.**



**MEAT
AND OFFAL
66 DAYS**



PRESENTATIONS



Solution
Injectable (Subcutaneous)
50 mL Vial
Box of 1 Vial



Solution
Injectable (Subcutaneous)
100 mL Vial
Box of 1 Vial



Solution
Injectable (Subcutaneous)
250 mL Vial
Box of 1 Vial

