



PASTEVAC BOVIN

***VACCINATE AGAINST
BOVINE PASTEURELLOSIS...***



***...WITH AN EFFECTIVE
AND EASY-TO-USE
INACTIVATED VACCINE***





RUMINANTS

VACCINES

PASTEovac BOVIN



INDICATIONS

Prevention



Bacteria

Inactivated vaccine against bovine pasteurellosis (*Pasteurella multocida*)



COMPOSITION

Vial Contains 50 Vaccine Doses

One Vaccine Dose (2 mL) Contains

- **PASTEURELLA MULTOCIDA SEROTYPE B** strain — 20×10^9 bacterial cells per dose
- Excipientsq.s. to 2 mL



ADMINISTRATION



PHARMACEUTICAL FORM

Injectable Solution

Injectable
Route



Subcutaneous
In the Neck



DOSAGE

Dilution

Use the product undiluted

Administration

Dose: 2 mL per animal
Only healthy animals should be vaccinated.

Duration of Use
Duration of Action



Single Dose
Booster: Every 6 months

IMMUNITY : Immunity develops 10 days after primary vaccination and lasts for 6 months.



WITHDRAWAL PERIODS



MEAT AND OFFAL
0 DAY



MILK
0 DAY



PRESENTATIONS



X1

- Solution
- Injectable Subcutaneous
- 100 mL Vial (50 Doses)

STORAGE

- 6 months at +20°C
- 1 year at +4°C

