

ANTI-INFECTIVES

SULFALOBS 33

RUMINANTS
SWINE - EQUINES
POULTRY - RABBITS



**TREAT COCCIDIOSIS AND
INFECTIONS CAUSED BY
BACTERIA SUSCEPTIBLE TO
SULFADIMERAZINE...**



**...WITH A VERSATILE
MULTI-SPECIES SOLUTION**



RUMINANTS
SWINE - EQUINES
POULTRY - RABBITS

SULFALOBS 33



INDICATIONS

Treatment



Coccidia



Bacteria

Bacteria susceptible to sulfadimerazine



COMPOSITION

100 mL of solution

SULFADIMERAZINE (as sodium salt) 33g

Excipients q.s. to 100mL



ADMINISTRATION



PHARMACEUTICAL FORM


Injectable
Route



Intramuscular
Intravenous



Injectable Solution



Oral
Route



Drinking Water



Oral Solution



DOSAGE



KG



3mL per 10kg BW

Duration of Use



**Divided into 2 injections per day
For 3 to 5 days**



5 mL per 1 L of drinking water

Duration of Use



For 3 to 5 days



WITHDRAWAL PERIODS



**MILK
5 DAYS**



**MEAT AND OFFAL
10 DAYS
POULTRY: 14 Days**



**EGGS
12 DAYS**



PRESENTATIONS



X1



Solution



Injectable or



100 mL Vial



Box of 1 Vial



X1



Solution



Injectable or



500 mL Vial

