



SYMPTEVAC

***VACCINATE RUMINANTS
AGAINST BLACKLEG...***



***...WITH AN EFFECTIVE
AND EASY-TO-USE
INACTIVATED VACCINE***





RUMINANTS

VACCINES

SYMPTEVAC



INDICATIONS

Prevention



Bacteria

*Inactivated vaccine against blackleg (*Clostridium chauvoei*)*



COMPOSITION

Vial Contains 50 or 100 Vaccine Doses
One Vaccine Dose

- **CLOSTRIDIUM CHAUVOEI ANACULTURE** — 10^9 bacterial cells per dose.
- Excipientsq.s. to 1 dose



ADMINISTRATION



PHARMACEUTICAL FORM

Injectable Solution

Injectable
Route



Subcutaneous
In the Neck



DOSAGE

Dilution

Use the product undiluted

Administration

Dose: 2 mL per bovine animal
Only healthy animals should be vaccinated.

Duration of Use
Duration of Action



Primary Vaccination

(One injection at 3–6 months of age,
followed by a second injection
3–4 weeks later)

Booster: Annually

IMMUNITY: develops 10 days after primary vaccination and lasts for 12 months.



WITHDRAWAL PERIODS



MEAT AND OFFAL
0 DAY



MILK
0 DAY



PRESENTATIONS



X1

- Solution
- Injectable – Subcutaneous
- 100 mL Vial 50 Doses

STORAGE

- 1 year at +20°C
- 2 years at +4°C

