

ANTI-INFECTIVES

RUMINANTS
POULTRY



TYL-DOX

**TREAT INFECTIONS CAUSED BY
BACTERIA SUSCEPTIBLE TO
TYLOSIN AND DOXYCYCLINE...**



**...WITH AN EFFECTIVELY
DOSED POWDER COMBINING
TWO SYNERGISTIC ANTIBIOTICS**





RUMINANTS
POULTRY

ANTI-INFECTIVES

TYL-DOX



INDICATIONS

Treatment



Bacteria

Bacteria susceptible to tylosin and doxycycline



COMPOSITION

100g of powder 

TYLOSIN (as tartrate) 10g

DOXYCYCLINE (as hyclate) 20g

Excipients q.s. to 100g



ADMINISTRATION



PHARMACEUTICAL FORM

Powder for Oral Solution

Oral Route



Drinking Water



or



DOSAGE



10 g per 20–40 L of drinking water



1 g per 40 kg BW

Duration of Use



For 3 to 5 days



WITHDRAWAL PERIODS



**MILK
NOT APPLICABLE**



**EGGS
NOT APPLICABLE**



MEAT AND OFFAL
Poultry: 7 DAYS
Calves: 12 DAYS



PRESENTATIONS



-  Powder
-  Drinking Water or Feed
-  Packet of 15g
-  Box of 100 packets



-  Powder
-  Drinking Water or Feed
-  Packet of 100g
-  Box of 25 packets



-  Powder
-  Drinking Water or Feed
-  1 kg Jar

